

Role of Particle Interactions in the Feshbach Conversion of Fermion Atoms to Bosonic Molecules

Jie Liu,^{1,2} Li-Bin Fu,¹ Bin Liu,¹ and Biao Wu³

¹*Institute of Applied Physics and Computational Mathematics, Beijing, 100088, China*

²*Center for Applied Physics and Technology, Peking University, 100084, Beijing, P.R.China*

³*Institute of Physics, Chinese Academy of Sciences, P.O.Box 603, Beijing 100080, China*

(Dated: February 1, 2008)

We investigate the Feshbach conversion of fermion atomic pairs to condensed boson molecules with a microscopic model that accounts the repulsive interactions among all the particles involved. We find that the conversion efficiency is enhanced by the interaction between boson molecules while suppressed by the interactions between fermion atoms and between atom and molecule. In certain cases, the combined effect of these interactions leads to a ceiling of less than 100% on the conversion efficiency even in the adiabatic limit. Our model predicts a non-monotonic dependence of the efficiency on mean atomic density. Our theory agrees well with recent experiments on ⁶Li and ⁴⁰K.

PACS numbers: 03.75.Ss, 05.30.Fk, 05.30.Jp, 03.75.Mn

Feshbach resonance has now become a focal point of the research activities in cold atom physics[1, 2, 3, 4] since its first experimental realization[5]. Among these research activities, the production of diatomic molecules from Fermi atoms with Feshbach resonance is of special interest and has attracted great attention. First, it is an interesting phenomenon by itself; second, it provides a unique experimental access to the BCS-BEC crossover physics[6]. So far, by slowly sweeping the magnetic field through the Feshbach resonance, samples of over 10^5 weakly bound molecules (binding energy ~ 10 kHz) at temperatures of a few tens of nK have been produced from quantum degenerate Fermi gas[7, 8, 9].

The Feshbach conversion is a complicated process involving many fermion atoms and boson molecules in a sweeping magnetic field that crosses a resonance. The theoretical description of the conversion efficiency as a function of sweep rate, atom mass, atomic density, and temperature is still under development. The existing theories include the Landau-Zener (LZ) model of two-body molecular production[10, 11] and its many-body extension at zero temperature[12, 13, 14], phase-space density model[15], and equilibration model at finite temperatures[16].

In this Letter we study a microscopic model of the Feshbach conversion that accounts all the two-body interactions, which include atom-atom, molecule-molecule, and atom-molecule interactions. These interactions are ignored in previous theoretical studies[11, 12, 13, 14]. We find that these interactions affect strongly the Feshbach conversion efficiency: the repulsive interaction between molecules tends to enhance the conversion efficiency while the other two repulsive interactions between atoms and between atom and molecule suppress the efficiency. Combined together, these interactions can yield a ceiling of less than 100% for the conversion efficiency even in the adiabatic limit of the sweeping magnetic field. This interaction-suppressed conversion efficiency is in spirit the same as the broken adiabaticity by interaction in the nonlinear LZ tunneling[17, 18]. In addition, our model

predicts a non-monotonic dependence of the conversion efficiency on mean atomic density. Our results are compared to recent experiments with ⁶Li and ⁴⁰K[8, 15]; they are in good agreement.

To include all particle interactions, we extend the two channel model[19, 20, 21] and write the Hamiltonian as

$$\begin{aligned}
 H = & \sum_{\mathbf{k}, \sigma} \epsilon_{\mathbf{k}} a_{\mathbf{k}, \sigma}^{\dagger} a_{\mathbf{k}, \sigma} + \left(\gamma + \frac{\epsilon_b}{2} \right) b^{\dagger} b \\
 & + \frac{U_a}{V_a} \sum_{\mathbf{k}, \mathbf{k}'} a_{\mathbf{k}, \uparrow}^{\dagger} a_{-\mathbf{k}, \downarrow}^{\dagger} a_{-\mathbf{k}', \downarrow} a_{\mathbf{k}', \uparrow} \\
 & + \frac{U_{ab}}{V_a} \sum_{\mathbf{k}, \sigma} a_{\mathbf{k}, \sigma}^{\dagger} a_{\mathbf{k}, \sigma} b^{\dagger} b + \frac{U_b}{V_b} b^{\dagger} b^{\dagger} b b \\
 & + \frac{g V_b}{V_a^{3/2}} \sum_{\mathbf{k}} \left(b^{\dagger} a_{-\mathbf{k}, \downarrow} a_{\mathbf{k}, \uparrow} + a_{\mathbf{k}, \uparrow}^{\dagger} a_{-\mathbf{k}, \downarrow}^{\dagger} b \right). \quad (1)
 \end{aligned}$$

Here $\epsilon_{\mathbf{k}} = \hbar^2 k^2 / 2m_a$ is the kinetic energy of the atom and $\sigma = \uparrow, \downarrow$ denote the two hyperfine states of the atom. $U_a = \Lambda U_0$, $U_{ab} = \Lambda U_1$, $g = \Lambda g_0$, and $\gamma = \gamma_0 - \Lambda g_0^2 / U_c$, where $\gamma_0 = \mu_{co}(B - B_0)$ is the molecule energy under the linearly changing magnetic field with $B = -\alpha_r t$, $g_0 = \sqrt{4\pi \hbar^2 a_{bg} \Delta B \mu_{co} / m_a}$ is the atom-molecule coupling [3], $U_0 = 4\pi \hbar^2 a_{bg} / m_a$ is the interaction between atoms, $U_1 = 4\pi \hbar^2 1.2 a_{bg} / m_{ab}$ is the atom-molecule scattering interaction, and $U_b = 4\pi \hbar^2 0.6 a_{bg} / m_b$ is the interaction between molecules[22]. $\Lambda \equiv (1 + U_0 / U_c)^{-1}$ and $U_c^{-1} = -\sum_{\mathbf{k}} (\exp(-k^2 / k_c^2) / 2\epsilon_{\mathbf{k}})$ with the cutoff momentum k_c representing the inverse range of interaction [21, 23, 24]. B_0 and ΔB are Feshbach resonance point and width, respectively. m_a and $m_b = 2m_a$ are masses for atoms and molecules, and $m_{ab} = \frac{2}{3}m_a$ is the reduced mass for the atom-molecule interaction.

In experiments, the molecular bosons are more tightly confined in space than the fermion atoms due to their different statistics. To reflect this, we use V_a for the volume of fermion atoms and V_b for bosonic molecules. We assume the zero temperature limit, consider only one

bosonic mode, and ignore all possible dissipations in the system, such as the loss of atoms by three-body collision.

In current experiments, the intrinsic energy width of a Feshbach resonance is much larger than the Fermi energy E_F [25], it is therefore reasonable to assume $\epsilon_{\mathbf{k}} = \epsilon$. This is called degenerate model in Ref.[12, 13, 20]. We introduce the following

$$\text{operators [12, 13], } L_x = \frac{\sum_{\mathbf{k}} (a_{\mathbf{k},\uparrow}^\dagger a_{-\mathbf{k},\downarrow}^\dagger b + b^\dagger a_{-\mathbf{k},\downarrow} a_{\mathbf{k},\uparrow})}{(N/2)^{3/2}}, L_y = \frac{\sum_{\mathbf{k}} (a_{\mathbf{k},\uparrow}^\dagger a_{-\mathbf{k},\downarrow}^\dagger b - b^\dagger a_{-\mathbf{k},\downarrow} a_{\mathbf{k},\uparrow})}{i(N/2)^{3/2}}, L_z = \frac{\sum_{\mathbf{k},\sigma} a_{\mathbf{k},\sigma}^\dagger a_{\mathbf{k},\sigma} - 2b^\dagger b}{N},$$

where $N = 2b^\dagger b + \sum_{\mathbf{k},\sigma} a_{\mathbf{k},\sigma}^\dagger a_{\mathbf{k},\sigma}$ is the total number of atoms. The Hamiltonian becomes

$$H = \frac{N}{4} \left(2\epsilon - \left(\gamma + \frac{\epsilon_b}{2} \right) - \frac{NU_a}{2V_a} - \frac{NU_{ab}}{V_a} \right) L_z - \frac{N^2}{16} \left(\frac{U_a}{V_a} + \frac{2U_{ab}}{V_a} - \frac{U_b}{V_b} \right) (1 - L_z)^2 + \frac{gV_b}{V_a^{3/2}} \left(\frac{N}{2} \right)^{3/2} L_x \quad [26].$$

With the commutators $[L_z, L_x] = \frac{4i}{N} L_y$, $[L_z, L_y] = -\frac{4i}{N} L_x$, $[L_x, L_y] = \frac{i}{N} (1 - L_z)(1 + 3L_z) + o(1/N^2)$, we can obtain the Heisenberg equations for the system $i\hbar \frac{d}{dt} L_l = [L_l, H]$, ($l = x, y, z$). Since all the commutators vanish in the limit of $N \rightarrow \infty$ and N is large in current experiments, it is appropriate to take L_x , L_y , and L_z as three real numbers u, v, w , respectively. These Heisenberg equations are then reduced to

$$du/d\tau = -\delta v - 2\chi v(1 - w), \quad dw/d\tau = \sqrt{2}v, \quad (2)$$

$$dv/d\tau = \frac{3\sqrt{2}}{4}(w - 1)(w + \frac{1}{3}) + \delta u + 2\chi u(1 - w), \quad (3)$$

where $\tau = (gV_b\sqrt{N}/\hbar V_a^{3/2})t$. Because of the identity $u^2 + v^2 = \frac{1}{2}(w - 1)^2(w + 1)$, with introducing the canonical variable $\theta = \arctan(v/u)$ we have a classical Hamiltonian,

$$\mathcal{H} = \delta w - \chi(1 - w)^2 + \sqrt{(w - 1)^2(w + 1)} \cos \theta. \quad (4)$$

The above equations show that all the experimental parameters affect the system via only two dimensionless parameters δ and χ . By a trivial shift of time origin, we can set $\delta = \alpha\tau$ with

$$\frac{\alpha_r}{\alpha} = \frac{4\pi\hbar n a_{bg} \Delta B}{m_a} \Lambda^2 \frac{V_b^2}{V_a^2}, \quad (5)$$

where $n = N/V_a$ is the mean atomic density. The nonlinear parameter χ is given by

$$\chi = \left(2.3 - \frac{0.15V_a}{\Lambda V_b} \right) \frac{V_a}{V_b} \sqrt{\frac{\pi\hbar^2 a_{bg} n}{m_a \mu_{co} \Delta B}}. \quad (6)$$

To understand the dynamics, we first look at the fixed points $\dot{w} = \dot{u} = \dot{v} = 0$ of Eqs.(2,3). The energies for these fixed points make up energy levels of the system as shown in Fig.1. One sees that the structure of these energy levels changes dramatically as the nonlinear parameter χ increases. Specifically, we observe: (i) There are two fixed points when $|\delta|$ is large enough: one for bosonic molecule (BM) and the other for fermion atom (FA). (ii) When $|\delta| < \delta_c = \sqrt{2}$, there is an additional fixed point




FIG. 1: Adiabatic energy levels for different interaction strengths. (a) $\chi = 0$; (b) $\chi = \chi_c = \sqrt{2}/4$; (c) $\chi = 1.5$. The unstable states are indicated by dashed lines (MQ and DM).

with $w = 1$. However, this fixed point is dynamically unstable[18]. (iii) For $\chi > \chi_c = \sqrt{2}/4$, there appears one more fixed point denoted by P_3 and, consequently, a loop in the energy levels. As we shall see, this loop has highly non-trivial physical consequences. This fixed point P_3 is also unstable.

Consider the adiabatic evolution of the system starting from a high negative value of δ with $w = 1$. This corresponds to the experiments where the magnetic field sweeps slowly across the Feshbach resonance with no bosonic molecules initially. When χ is small, such as in Fig.1(a), the evolution of the system follows the solid line, converting all fermion atoms into molecules. However, when χ is beyond χ_c as in Fig.1(c), the system will find no stable energy level to follow at singular point M . As a result, only a fraction of fermion atoms are converted into bosonic molecules.

This simple analysis is confirmed by our numerical results, which are plotted in Fig.2. In our calculation, the 4-5th Runge-Kutta step-adaptive algorithm is used in solving the differential equations (2,3). Because $w = 1$ is a fixed point when $\delta < -\sqrt{2}$, we start from $(w, u, v) \approx (1, 0, 0)$ and sweep the field from $\delta = -\sqrt{2}$ to 200. In Fig.2, the conversion efficiency Γ , i.e., the fraction of the converted fermion atom pairs is drawn as a function of α . Evidently, Γ approaches one as $\alpha \rightarrow 0$ when $\chi < \chi_c$, indicating that all atomic pairs are converted into molecules. In contrast, when $\chi > \chi_c$, Γ does not increase to one in the adiabatic limit $\alpha \rightarrow 0$. This means that there is a ceiling $\Gamma_{ad} (< 100\%)$ on the conversion efficiency. Moreover, Fig.2 demonstrates that positive χ suppresses the conversion efficiency whereas the negative χ enhances it. Because the repulsive interaction between bosonic molecules enters χ as a negative value, it enhances the conversion efficiency; the repulsive fermion atomic interaction and atom-molecule interaction contribute positively to χ , they suppress the conversion.

The ceiling Γ_{ad} on the atom-molecule conversion ef-




FIG. 2: Conversion efficiency Γ as a function of the sweeping rate α for various interactions.

efficiency depends on χ . This dependence can be found by examining the phase space diagrams of our system shown in Fig.3. As δ ramps up slowly from a large negative value, the fixed point P_3 will move up until it hits the fixed point $w = 1, u = 0, v = 0$, represented by a dark straight line in Fig.3(a). This collision occurs at $\delta = -\sqrt{2}$. Immediately after the collision, the hyperbolic fixed point P_3 is no longer a fixed point and becomes a solution that evolves along the dark line in Fig.3(b). The dark line is given by $\sqrt{2} = \chi(1 - w) - \sqrt{1 + w \cos \theta}$, which is found by taking $E = \delta = -\sqrt{2}$ in the Hamiltonian (4). As the action of this trajectory is nonzero while a fixed point has zero action, this collision of the two fixed points represents a sudden jump in action. It is this sudden jump that has caused the nonzero fraction of remnant atoms. As δ ramps up further slowly, the trajectory will change its shape as witnessed in Fig.3(c); however, its action stays constant as demanded by the classical adiabatic theorem[27, 28]. The action is

$$I = \begin{cases} \frac{1}{2\pi} \oint \frac{\cos \theta \sqrt{8\chi^2 - 4\sqrt{2}\chi + \cos^2 \theta}}{2\chi^2} d\theta, & \frac{\sqrt{2}}{4} < \chi < \frac{\sqrt{2}}{2}; \\ \frac{1}{2\pi} \int_0^{2\pi} \frac{4\chi^2 - 2\sqrt{2}\chi + \cos^2 \theta}{2\chi^2} d\theta, & \chi > \frac{\sqrt{2}}{2}, \end{cases} \quad (7)$$

which yields the ceiling on the efficiency

$$\Gamma_{ad} = 1 - \frac{1}{2}I = \frac{4\sqrt{2}\chi - 1}{8\chi^2}, \chi > \frac{\sqrt{2}}{4}; \quad \Gamma_{ad} = 1, \chi < \frac{\sqrt{2}}{4}. \quad (8)$$

Now we compare our theory with existing experiments. For the experiment with ^6Li [8], the mean density is $n = 4 \times 10^{12} \text{cm}^{-3}$ with $N = 6 \times 10^5$ atoms. The scattering length $a_{bg} = 59a_B$, $\mu_{co} \sim 2\mu_B$, where a_B and μ_B are Bohr radius and Bohr magneton, respectively, and the resonance width $\Delta B = 0.1 \text{G}$ at $B_0 = 543.8 \text{G}$. The Fermi energy E_F in the combined harmonic and box-like trapping potential of Ref.[8] is given by $E_F = [15\pi\hbar^3\omega_r^2 / (8\sqrt{2}m_aL)]^{2/5}$, where $\omega_r = 2\pi \times 800 \text{s}^{-1}$ is the angular frequency of the radial harmonic trap and $L = 480 \mu\text{m}$ is the size of the axial po-




FIG. 3: Phase spaces of Hamiltonian (4). The dark line in (a) is for the fixed point $w = 1, u = 0, v = 0$. It is a line because θ is not defined at $u = v = 0$. The two fixed points on line $w = 1$ in (c) are in fact the same fixed point; they are artifact caused by the definition $\theta = \arctan(v/u)$.

tential. The ground state energy of molecular bosons is $E_G = \hbar\omega_r + \hbar^2\pi^2/(2m_bL^2)$. Then we have $V_a/V_b = E_F/E_G = 36$. We set $\Lambda = 391$ with a momentum cut-off $K_c = 96k_F$ [29]. From Eq.(5), the sweeping rate is $\alpha_r/\alpha = 20 \text{G/ms}$. The second term in the bracket of Eq.(6) that accounts for the repulsive interaction between bosonic molecules is small, so the interaction parameter takes the form of $\chi \simeq 2.3 \frac{V_a}{V_b} \sqrt{\frac{\pi\hbar^2 a_{bg} n}{m_a \mu_{co} \Delta B}}$. From the above experimental parameters, we find the interaction parameter as $\chi = 1.26$. This strong interaction ($> \chi_c$) indicates a ceiling of $\Gamma_{ad} = 0.48$ via Eq.(8). This is in good agreement with experiments (see Fig.4a).

For ^{40}K , the situation is different. The resonance at $B_0 = 202.1 \text{G}$ has a large width of $\Delta B = 7.8 \text{G}$ and the mass of ^{40}K is 7 times that of ^6Li . In Ref.[15], the fermions are confined in a dipole trap characterized by a radial frequencies ν_r between 312 and 630 Hz and an aspect ratio of $\nu_r/\nu_z = 70$. The Fermi energy is $E_F = \hbar(3N\omega_r^2\omega_z)^{1/3}$ and the ground state energy of condensed bosons is $E_G = \hbar\omega_r + \hbar\omega_z/2$. For the dipole trap, the ratio $V_a/V_b = (E_F/E_G)^{3/2} = 251$. With $a_{bg} = 174a_B$, $\mu_{co} \sim 2\mu_B$, initial clouds have mean densities $n = 2 \times 10^{12} \text{cm}^{-3}$, and $N = 2.5 \times 10^5$ [15], we obtain $\chi = 0.135 \sim 0.274$, which is less than the threshold $\chi_c = \sqrt{2}/4$. Therefore, ^{40}K atom pairs can be completely converted to bosonic molecules in adiabatic limit. Indeed, the conversion efficiency up to 90% has been observed[15].

We emphasize that the suppressed conversion efficiency by particle interaction dominates only at low temperatures. As a result, in the above we have only compared to the data obtained at low temperatures ($T/T_F = 0.1$ for ^6Li and $T/T_F = 0.05$ for ^{40}K). Temperature can affect the conversion efficiency strongly as reported in Ref.[15]. The ceiling of 50% conversion efficiency observed in Ref.[7] is likely a thermal effect since the experiment is performed at $T/T_F = 0.33$, and has been explained by the theories of finite temperature[30, 31]

For the ^6Li , from Eq.(2)(3) we have also calculated numerically the conversion efficiency as a function of sweeping rate. The comparison between our theory and exper-




FIG. 4: (a) Comparison between our theory and experimental data of ^6Li [8] for the conversion efficiency Γ as a function of field sweep rates. (b) The dependence of Γ on the mean atomic density.

iment is shown in Fig.4a. They are in a good agreement. In addition, our model predicts a non-monotonic depen-

dence of the conversion rate on the mean atomic density (see Fig.4(b)). This can be understood from Eqs.(5,6). In Eq.(5), we see the effective sweeping rate α is inversely proportional to the atomic density. So, increasing the density will reduce the effective sweeping rate and therefore enhance the conversion rate. On the other hand, higher density will give larger nonlinearity χ as indicated in Eq.(6), which in turn suppresses the atom-molecule conversion. These two factors compete with each other, giving rise to the non-monotonic curves in Fig.4(b). In practical experiments, to achieve higher conversion efficiency, one needs to carefully choose initial fermion atom density, making it fall into the optimal parameter regime.

In summary, we have identified the significant role of the interactions between particles in the Feshbach conversion of atomic fermion pairs to molecular bosons. Our theory is consistent with the existing experiments. Our model also predicts a non-monotonic dependence of the conversion rate on the mean atomic density, which is important for the optimal choice of parameters in future Feshbach experiments.

This work was supported by NSF of China (10725521,10604009,10504040) and the 973 project (2006CB921400,2007CB814800). B.W. is also supported by the “BaiRen” program of the CAS.

-
- [1] E. Timmermans, P. Tommasini, M. Hussein, and A. Kerman, Phys. Rep. **315**, 199 (1999).
 - [2] R.A. Duine and H.T.C. Stoof, Phys. Rep. **396**, 115 (2004).
 - [3] Q. Chen, J. Stajic, S.N. Tan, and K. Levin, Phys. Rep. **412**, 1 (2005).
 - [4] T. Köhler, K. Góral, and P.S. Julienne, Rev. Mod. Phys. **78**, 1311 (2006).
 - [5] S. Inouye *et al.*, Nature **392**, 151 (1998).
 - [6] C.A. Regal, M. Greiner, and D.S. Jin, Phys. Rev. Lett. **92**, 040403 (2004).
 - [7] C.A. Regal *et al.*, Nature (London) **424**, 47 (2003).
 - [8] K.E. Strecker, G.B. Partridge, and R.G. Hulet, Phys. Rev. Lett. **91**, 080406 (2003).
 - [9] J. Cubizolles *et al.*, Phys. Rev. Lett. **91**, 240401 (2003).
 - [10] C. Zener, Proc. R. Soc. A **137**, 696 (1932); L.D. Landau and E.M. Lifshitz, *Quantum Mechanics* (Pergamon, Oxford, 1977).
 - [11] K. Góral *et al.*, J. Phys. B **37**, 3457 (2004).
 - [12] E. Pazy *et al.*, Phys. Rev. Lett. **95**, 170403 (2005).
 - [13] I. Tikhonenkov *et al.*, Phys. Rev. A **73**, 043605 (2006).
 - [14] E. Altman and A. Vishwanath, Phys. Rev. Lett. **95**, 110404 (2005).
 - [15] E. Hodby *et al.*, Phys. Rev. Lett. **94**, 120402 (2005).
 - [16] J.E. Williams *et al.*, J. Phys. B, **37** (2004) L351-L357
 - [17] B. Wu and Q. Niu, Phys. Rev. A **61**, 023402 (2000).
 - [18] J. Liu *et al.*, Phys. Rev. A **66**, 023404 (2002).
 - [19] J. Javanainen *et al.*, Phys. Rev. Lett. **92**, 200402 (2004); R. A. Barankov and L. S. Levitov, *ibid.* **93**, 130403 (2004); A. V. Andreev *et al.*, *ibid.* **93**, 130402 (2004); J. Dukelsky *et al.*, *ibid.* **93**, 050403 (2004).
 - [20] T. Miyakawa and P. Meystre, Phys. Rev. A **71**, 033624 (2005).
 - [21] W. Yi and L.-M. Duan, Phys. Rev. A **73**, 063607 (2006).
 - [22] Here the a_{bg} of atom-molecule and molecule-molecule is taken to be 1.2 and 0.6 times that of atom-atom, respectively, see D. S. Petrov, C. Salomon, and G. V. Shlyapnikov, Phys. Rev. Lett. **93**, 090404 (2004).
 - [23] S. J. J. M. F. Kokkelmans *et al.*, Phys. Rev. A **65**, 053617 (2002).
 - [24] Q. Chen and K. Levin, Phys. Rev. Lett. **95**, 260406 (2005).
 - [25] R.B. Diener and T.-L. Ho, cond-mat/0405174.
 - [26] In deducing the atom-atom scattering term we need introduce the collective pseudo-spin operators $\hat{S}^+ = \sum_{\mathbf{k}} a_{\mathbf{k}\uparrow}^\dagger a_{-\mathbf{k}\downarrow}^\dagger$, $\hat{S}^- = \sum_{\mathbf{k}} a_{-\mathbf{k}\downarrow} a_{\mathbf{k}\uparrow}$, $\hat{S}_z = \sum_{\mathbf{k}} \frac{1}{2} (a_{\mathbf{k}\uparrow}^\dagger a_{\mathbf{k}\uparrow} + a_{-\mathbf{k}\downarrow}^\dagger a_{-\mathbf{k}\downarrow} - 1)$. It is easy to prove that $\hat{S}^2 = \hat{S}_z^2 - \hat{S}_z + \hat{S}^+ \hat{S}^-$ is a conservation and $S = N/4$. Combining the conserved relation of the total particles, $N/4 = \hat{b}^\dagger \hat{b} + \hat{S}_z$, we can rewrite the atom-atom scattering term as $\hat{S}^+ \hat{S}^- = \frac{1}{2} \sum_{\mathbf{k}, \sigma} a_{\mathbf{k}, \sigma}^\dagger a_{\mathbf{k}, \sigma} \hat{b}^\dagger \hat{b} + \frac{N}{2} - \hat{b}^\dagger \hat{b}$.
 - [27] L.D. Landau and E.M. Lifshitz, *Mechanics* (Pergamon, Oxford, 1977).
 - [28] J. Liu, B. Wu, and Q. Niu, Phys. Rev. Lett. **90**, 170404 (2003).
 - [29] The cutoff k_c is chosen such that the tunnelling window $2\delta_c$ for converting atomic fermions to molecular bosons is consistent with the Feshbach resonance width $\mu_{co}\Delta B$ of ^6Li .
 - [30] E. Pazy, A. Vardi, and Y. B. Band, Phys. Rev. Lett. **93**, 120409 (2004).
 - [31] J. Chwedeńczuk, K. Góral, T. Köhler, and P. S. Julienne,

Phys. Rev. Lett. **93**, 260403 (2004).